

Lecture 18

Taking a picture of a black hole:
computational imaging taken to an extreme



6.869/6.819 Advances in Computer Vision


Bill Freeman, Phillip Isola

spring 2022

April 7, 2022

The black hole story, told in the style of *Rosencrantz and Guildenstern are Dead*

Many slides courtesy of Event Horizon
Telescope Imaging Working Group, and
Katie Bouman



Rosencrantz and Guildenstern are Dead

Hamlet

POLONIUS: My Lord! I have news to tell you.

HAMLET (releasing ROS and mimicking): My lord, I have news to tell you... When Rodius was an actor in Rome...

(ROS comes down to re-join GUIL.)

POLONIUS (as he follows HAMLET out): The actors are come hither my lord.

HAMLET: Buzz, buzz.

(Exeunt HAMLET and POLONIUS.)

(ROS and GUIL ponder. Each reluctant to speak first.)

GUIL: Hm?

ROS: Yes?

GUIL: What?

ROS: I thought you...

GUIL: No.

ROS: Ah.

(Pause.)

GUIL: I think we can say we made some headway.

ROS: You think so?

GUIL: I think we can say that.

ROS: I think we can say he made us look ridiculous.

GUIL: We played it close to the chest of course.

ROS (derisively): "Question and answer. Old ways are the best ways"! He was scoring off us all down the line.

GUIL: He caught us on the wrong foot once or twice, perhaps, but I thought we gained some ground.

ROS (simply): He murdered us.

GUIL: He might have had the edge.

ROS (roused): Twenty-seven - three, and you think he might have had the edge?! He murdered us.



Shep Doleman



MIT's Haystack Observatory, Westborough, MA





Shep Doeleman <dole@haystack.mit.edu>

Mon, Sep 23, 2013, 7:37 PM

to Bill, Katie, Daniel, Rusen, Victor, Vincent ▾

Dear Bill, Dan and Katie,

It was great to talk with you all and to hear about Dan and Katie's interesting work - I don't think I'll look at leaves or fabric in the same way again.

Your ideas and gameplan sound reasonable, and is along the lines I was hoping for - a fresh look at this problem with unjaded astronomer eyes. There will undoubtedly be course corrections along the way, but this is a good start. We can generate fourier plane baseline tracks for you, or we can supply routines that will do that for you (functions of telescope location on the Earth, Greenwich mean time, and the sky position of the source).


I wonder to what extent the textured approach will contain the imprint of the types of images we expect to see (a 'gentle' prior), but let's see what happens. I really like the idea of generating a family of images that all obey the observations, but differ where we have no fourier data.

All the Best, Shep

<https://www.almaobservatory.org/en/press-releases/astronomers-capture-first-image-of-a-black-hole/>

In the shadow of a black hole

1:13 - 3:50 or 5:05.



Hubble Telescope


(optical wavelength)


NASA, ESA and the Hubble Heritage Team (STScI/AURA); Acknowledgment: P. Cote (Institute of Astrophysics) and E. University) Herzberg



This artist's impression depicts the black hole at the heart of the enormous elliptical galaxy Messier 87 (M87). This black hole was chosen as the object of paradigm-shifting observations by the Event Horizon Telescope. The superheated material surrounding the black hole is shown, as is the relativistic jet launched by M87's black hole.
Credit: ESO/M. Kornmesser

Black Hole Simulation





$$\theta = 1.22 \frac{\lambda}{D}$$

The Fourier Transform relationship between the light amplitudes entering the aperture of the telescope and those focussed at the sensor impose a resolution limit on the image from an optical system.

black hole photon
capture radius:

$$R_c = \frac{\sqrt{27}GM}{c^2}$$

measured wavelength

$$1.22 \frac{1.3mm}{12.7 \times 10^6 m} = 25.7 \mu\text{arc seconds}$$

diameter of Earth

Predicted black hole mass: 3-6 Billion solar masses,
55M light years away implying shadow size: between 42 and
20 micro arc seconds

Earth Sized


How Big Must Our Telescope Be?

Telescope
Size
13 million meters

Wavelength
 1.3 mm
Angular Resolution
 $20 \mu\text{as}$



Simulation of M87



Ideal Image with
Earth-Sized Telescope





Figure 2: The same baseline as in Figure 1, but for waves incident from an angle θ from the vertical. The waves arrive at the antennas again exactly in phase, because the angle is such that the difference in path length is λ .

“We run a double-slit interference experiment in reverse” —Shep Doleman

Double Slit Interference




VAN CITTERT-ZERNIKE THEOREM

The theorem states that, for ideal sensors, the time-averaged correlation of the measured signals from two telescopes, i and j , for a single wavelength, λ , can be approximated as:

$$\Gamma_{i,j}(u, v) \approx \int_{\ell} \int_m e^{-i2\pi(u\ell + vm)} I_{\lambda}(\ell, m) d\ell dm \quad (1)$$

where $I_{\lambda}(\ell, m)$ is the emission of wavelength λ traveling from the direction $\hat{s} = (\ell, m, \sqrt{1 - \ell^2 - m^2})$. The dimensionless coordinates (u, v) (measured in wavelengths) are the projected baseline, B , orthogonal to the line of sight.¹

The Event Horizon Telescope (EHT)



Event Horizon Telescope (EHT)

A Global Network of Radio Telescopes


2018 Observatories






Imaging a Black Hole with the Event Horizon Telescope

EHT Collaboration



FOURIER DOMAIN COVERAGE OF THE 4 NIGHTS OF OBSERVATIONS




Fourier domain

Because the image is real valued, the Fourier transform is Hermitian.
K telescopes gives K chose 2 baselines, or observed Fourier frequencies.
As the Earth rotates over a night, the projected baselines sweep out elliptical paths in Fourier space.

ICCV 2015 was in Santiago, Chile

Shep said we should go see the Alma radio telescopes in Atacama, and I arranged for a trip there, with Bernhard Scholkopf, Yoav Schechner, and Katie Bouman.



The Atacama Telescope Project

HomeGoalsScienceSite InfoSite SurveysWorkshops

THE ATACAMA REGION

The site under consideration for the Atacama Telescope is located in the Chajnantor Region of Northern Chile, in the vicinity of the site chosen for the ALMA (formerly MMA) radio array, to be built by a consortium between the National Radio Astronomy Observatory and the European Southern Observatory.

Recently, attention has focused on [Cerro Negro](#) and [Cerro Chajnantor](#) as a potential sites for the Atacama telescope.


The Atacama Desert extends over northern Chile. The Chajnantor Plateau is located about 20 km from the Bolivian border, and some 80 km from the Argentine border, about 1500 km north of Santiago. The Salar de Atacama, a salt flat some 90 km long and 40 km wide, is 40 km to the west. The nearest inhabited village is San Pedro de Atacama (pop. 1200), at the northern tip of the Salar. The nearest airport with regular commercial service is in Calama (pop. 120,000), 150 km to the west. Calama is also a major center of the copper mining industry. The regional capital of Antofagasta (pop. 250,000), on the coast, is more than 300 km to the west of Chajnantor.

The region is likely to become the center of a major concentration of astronomical observatories,

with the highest sites in the world. The surrounding peaks range in altitude between 5,000 and 6,000 m (17,000 and 20,000 ft). The combination of geographical location and altitude also makes the region among the driest on Earth.

Since 1998, [Tests](#) on a 150 m rise above the ALMA site have shown very low values for the median optical seeing and for the perceptible water vapor.

In June of 1996, the then President of Chile, don Eduardo Frei, signed a bill naming the Chajnantor region a "National Science Preserve". The land concession was transferred to the Chilean National Committee for Science and Technology (CONICYT). Representatives of AUI (Associated Universities, Inc.), NRAO and Cornell were invited to the ceremony.



Map of Chile from the [CIA World Factbook](#). The dashed square represents the Upper Atacama Desert Region discussed here. The ALMA site (filled red circle), the city of Calama (open blue circle), and the town of San Pedro de Atacama (filled blue circle) are indicated.













Timelapse videos from Bernhard, from Atacama, 2015



atacama milkyway

225 views • 3 years ago

<https://youtu.be/O86mycdxtYM?t=12>



atacama tree

111 views • 3 years ago

<https://youtu.be/8YOqdAooO9A>

The Atacama Large Millimeter/submillimeter Array (ALMA) by night, u...



Eso Observatory

MAR 2013






Dec. 12, 2015
Below ALMA
telescope array,
Atacama, Chile
9,000 feet

Yoav in pressurized car,
going to 15,000 feet



ALMA telescope array, 15,000 feet. Atacama, Chile, 2015





ALMA telescope array, 15,000 feet. Atacama, Chile, 2015



ALMA telescope array, 15,000 feet. Atacama, Chile, 2015







Astronomical Radio Imaging – Centaurus A Galaxy





Very Long Baseline Interferometry (VLBI)




The Computational Imaging Problem

True Image



Sparse
Measurements



Reconstruction





ALGORITHM




The Computational Imaging Problem

True Image




Sparse
Measurements




Reconstruction



Inverse Fourier Transform



Inverting the Imaging System: Ambiguity

















Figure 10: *a)* An example (model) sky map. *d)* The corresponding visibilities (Fourier Transform of the map). *e)* The synthesized beam, or point-spread-function, of a model antenna array. *e)* The sampling function of the array, whose Fourier Transform gives the beam in (b). *f)* The product of panels (d) and (e), representing the sampled visibilities. These are the actual measurements from the array. *c)* The dirty map that results from the Fourier Transform of the sampled visibilities. This is the same as the convolution of the map in (a) and the synthesized beam in (b).



Katie presenting EHT poster at CVPR 2016

	BLACK HOLE	
TARGET		
CLEAN		
SQUEEZE		
BSMEM		
CHIRP		

Some of the image priors explored for
Event Horizon Telescope black hole image
reconstructions one has ever seen
(what assumptions do you make about an image of something that
no one has ever seen before?)

- Positivity: the light intensity must be positive.
- Compactness: The source has a finite size
- Image entropy
- Image smoothness
- Image sparsity in the pixel, or gradient, domains

VLBI Reconstruction Dataset

A Dataset Designed to Train and Test Very Long Baseline Interferometry Image Reconstruction Algorithms

HOME FAQ TRAINING DATA REAL DATA TEST DATA SCOREBOARD RESULT GALLERY GENERATE YOUR DATA EHT IMAGING CHALLENGE

EHT Imaging Challenge

Welcome to the Event Horizon Telescope Imaging Challenge Webpage! This challenge is meant to help us understand the performance of different imaging algorithms on future Event Horizon Telescope (EHT) data. We hope the results of the challenge will help us better understand the biases of each imaging algorithm, and aid in developing better methods.

Next Deadline: December 20, 2017

- [Testing Data and Submission Instructions](#)
- [Data Parameters and Noise Properties](#)
- [Sample Data With Ground Truth Images](#)
- [Past Challenges](#)
- [Data Formats and Conversion](#)
- [Sample Imaging Script](#)
- [Questions and Feedback](#)

Testing Data and Submission Instructions

1. **Download the test data from [HERE](#).**
2. **Use your algorithm to generate an image for each of the data files.** For each < filename >.txt file, submit a FITS image with the name < filename >.fits and the FOV specified in the README file. Further instructions can be found in the README file.
3. **Submit your reconstructed images.** Compress all of your reconstructed FITS images into a ZIP file. Submit this ZIP file with the required additional information.

Method Name: Email: Images:

Additional Information (such as website/code links):

Sample Data With Ground Truth Images


We provide a set of sample data, along with their ground truth images, to help in getting your imaging algorithms working on the blind, test data.

Static Emission

You can download the sample data from [here](#). This sample data was generated with the same telescope parameters as the blind, test data. We have included data without any systematic errors or atmospheric errors, data with just atmospheric errors, and data with both systematic and atmospheric errors. Their naming is as follows:

Filename	Property
challenge_x_wNoPhaseError	Only thermal noise included in visibility measurements
challenge_x	Thermal and phase (atmospheric) errors included in visibility measurements
challenge_x_wSystematics	Thermal, amplitude (systematic) and phase (atmospheric) errors included in visibility measurements

Sample Ground Truth Images



Sample Number	Source Location	Telescopes	Total Flux (Janskys)	Field of View (arcseconds)
1	SgrA*	ALMA, SMT, LMT, SMA, SPT, CARMA, PV, PdBI, KP, Haystack	2	0.00016
2	SgrA*	ALMA, SMT, LMT, SMA, SPT, CARMA, PV, PdBI, KP, Haystack	2	0.00025
3	SgrA*	ALMA, SMT, LMT, SMA, PV, SPT, KP, PdBI	2	0.00016
4	SgrA*	ALMA, SMT, LMT, SMA, PV, SPT	2	0.00016
5	M87	ALMA, SMT, LMT, SMA, SPT, CARMA, PV, PdBI, KP, Haystack	2	0.00010
6	M87	ALMA, SMT, LMT, SMA, SPT, CARMA, PV, PdBI, KP, Haystack	2	0.00010
7	M87	ALMA, SMT, LMT, SMA, PV, SPT, KP, PdBI	2	0.00025
8	M87	ALMA, SMT, LMT, SMA, PV	2	0.00010

Extreme Imaging via Physical Model Inversion: Seeing Around Corners and Imaging Black Holes

by

Katherine L. Bouman

B.S.E., Electrical Engineering, University of Michigan, 2011

S.M., Electrical Engineering and Computer Science, M.I.T., 2013

Submitted to the Department of Electrical Engineering and Computer Science
in partial fulfillment of the requirements for the degree of

Doctor of Philosophy
in Electrical Engineering and Computer Science
at the Massachusetts Institute of Technology

September 2017

© 2017 Massachusetts Institute of Technology
All Rights Reserved.

Signature of Author: _____

Department of Electrical Engineering and Computer Science
August 31, 2017

Certified by: _____

Professor William T. Freeman
Thomas and Gerd Perkins Professor of Electrical Engineering and Computer Science
Thesis Supervisor

Katie and EHT team members at radio telescope in Mexico, April 2, 2017



Observations April 5-11, 2017

All sites were technically ready and with good weather on the first night of the observing window. Observations were triggered on 2017 April 5, 6, 7, 10, and 11. Table 1 shows the median zenith sky opacities for each of the triggered days. April 8 was not triggered due to thunderstorms at the LMT, SMT shutdown due to strong winds, and the need to run technical tests at ALMA. April 9 was not triggered due to a chance of the SMT remaining closed due to strong winds and LMT snow forecast. [Weather was good to excellent for all other stations throughout the observing window.](#)

...

Observations from the EHT's 2017 April campaign are the first ever to have the necessary sensitivity, coverage, and resolution for horizon-scale imaging of black hole candidates M87 and SgrA*.

2nd way the EHT got lucky

For M87, the expected shadow diameter is 19–38 μ as.


....

We present the first Event Horizon Telescope (EHT) images of M87, using observations from April 2017 at 1.3 mm wavelength. These images show a prominent ring with a diameter of $\sim 40 \mu$ as,





Calibration










The correlator is a special-purpose supercomputer that combines data from the telescopes, to recover measurements that would be seen from an Earth-size telescope.

Calibration algorithms find the weak signals hiding in the correlator output, and more precisely tune the data to extract a stronger signal.










The Event Horizon Telescope






The Event Horizon Telescope






Lo-band eht-imaging on April 11: slowly building up data



Team 1 – End of the First Day of Imaging M87



(mention my contribution)



EHT Imaging Working Group



The dangers of false confidence and collective confirmation bias are magnified for the EHT because the array has fewer sites than typical VLBI arrays, there are no previous VLBI images of any source at 1.3 mm wavelength, and there are no comparable black hole images on event-horizon scales at any wavelength.

We subdivided our first M87 imaging efforts into four separate imaging teams. The teams were blind to each others' work, prohibited from discussing their imaging results and even from discussing aspects of the data that might influence imaging (e.g., which stations or data might be of poor quality).



Imaging Stage 1/2: Blind Imaging Comparisons

Team 1

Region:
The Americas
(SAO, UoA, U.Concepcion)


The Imaging WG was divided
into four independent teams

Team 4

Region:
East Asia (ASIAA, KASI,
NAOJ)

Team 2

Region:
Global
(MIT Haystack, Radboud U, NAOJ)



Team 3

Region:
Cross-Atlantic
(MPIfR, Boston U, IAA,
Aalto)

Each team blindly
reconstructed images
Goal: Assess human bias



The First EHT Images of M87

July 24, 2018

Team 1

Region:
The Americas
(SAO, UoA, U.Concepcion)

Team 4


Region:
East Asia (ASIAA, KASI,
NAOJ)

Team 2


Region:
Global
(MIT Haystack, Radboud U, NAOJ)

Team 3

Region:
Cross-Atlantic
(MPIfR, Boston U, IAA, Aalto)



Each team blindly
reconstructed images
Goal: Assess human bias



[Zoom In](#) [Zoom Out](#) [Reset image size](#)


Figure 4. The first EHT images of M87, blindly reconstructed by four independent imaging teams using an early, engineering release of data from the April 11 observations. These images all used a single polarization (LCP) rather than Stokes I , which is used in the remainder of this Letter. Images from Teams 1 and 2 used RML methods (no restoring beam); images from Teams 3 and 4 used CLEAN (restored with a circular $20 \mu\text{as}$ beam, shown in the lower right). The images all show similar morphology, although the reconstructions show significant differences in brightness temperature because of different assumptions regarding the total compact flux density (see Table 2) and because restoring beams are applied only to CLEAN images.



The Event Horizon Telescope

The First EHT Images of M87


July 24, 2018



2nd EHT Imaging Workshop



Imaging Stage 2/2: Imaging Parameter Survey



Imaging algorithms were tested on a suite of synthetic datasets

Goal: Optimize imaging algorithms with objective performance assessment

Imaging Stage 2/2: Imaging Parameter Survey







Figure 7. Selection of the `eht-imaging` (RML) parameter survey results on real and synthetic data with April 11 EHT baseline coverage. A 2D slice of the 7D parameter space is displayed, corresponding to different weights on the MEM and TV regularizers. All other parameters are kept constant (Compact Flux = 0.6 Jy, Initial/MEM FWHM = $40\ \mu\text{as}$, Systematic Error = 1%, TSV = 0, and $\ell_1 = 0$). The left panel shows results of the parameter search on the Crescent synthetic data, while the right panel shows reconstructions for the same parameters on M87 data. Images that meet the threshold for the Top Set are outlined in green. Note that the upper-left reconstruction has *no* regularization; it is produced by enforcing only image positivity and a constrained FOV.







Figure 10. Cross-validation of the imaging parameter selection procedure. In each of the left four columns, we show reconstructed images for the simple geometric source models. These reconstructions do not use the fiducial imaging parameters identified by the full training set; instead, we selected the imaging parameters for each geometric source model after excluding that particular model from the parameter selection process. For example, in the disk reconstructions, the parameters were selected by assessing reconstructions of only the ring, crescent, and double source models. Thus, the selected parameters vary among these four columns, but we can verify that the training sets do not overly constrain the outcomes. In the fifth column, we show reconstructions of a GRMHD snapshot (Paper [V](#)) using the fiducial M87 parameters selected from all four geometric models. That is, the script and parameters used to produce these GRMHD image reconstructions are identical to those used to produce our fiducial M87 images (shown in Figure [11](#)). Because the GRMHD snapshot has a substantially higher peak brightness than the reconstructions, its column has been scaled to the peak brightness of the `eht-imaging` reconstruction.




Fiducial images for all four days and three scripts

- Best images out of 1008, 37500, and 10800 images surveyed by the Difmap, eht-library, and SMILI scripts, respectively
- All images from the four different observing days show the asymmetric ring structure corresponding to the black hole shadow






**Fiducial images of M87 for April 11 restored
to an equivalent resolution show
remarkably similar structure**



Comparison of reconstructions, in the Fourier domain



Focus on the First Event Horizon Telescope Results

Shep Doeleman (EHT Director) on behalf of the EHT Collaboration

April 2019




Figure 1. EHT images of M87 on four different observing nights. In each panel, the white circle shows the resolution of the EHT. All four images are dominated by a bright ring with enhanced emission in the south. From Paper IV (Figure 15).

We report the first image of a black hole.

This Focus Issue shows ultra-high angular resolution images of radio emission from the supermassive black hole believed to lie at the heart of galaxy M87 (Figure 1). A defining feature of the images is an irregular but clear bright ring, whose size and shape agree closely with the expected lensed photon orbit of a 6.5 billion solar mass black hole. Soon after Einstein introduced general relativity, theorists derived the full analytic form of the photon orbit, and first simulated its lensed appearance in the 1970s. By the 2000s, it was possible to sketch the "shadow" formed in the image when synchrotron emission from an optically thin accretion flow is lensed in the black hole's gravity. During this time, observational evidence began to build for the existence of black holes at the centers of active galaxies, and in our own Milky Way. In particular, a steady progression in radio astronomy enabled very long baseline interferometry (VLBI) observations at ever-shorter wavelengths, targeting supermassive black holes with the largest apparent event horizons: M87, and Sgr A* in the Galactic Center. The compact sizes of these two sources were confirmed by studies at 1.3mm, first exploiting baselines that ran from Hawai'i to the mainland US, then with increased resolution on baselines to Spain and Chile.




Figure 2. A map of the EHT. Stations active in 2017 and 2018 are shown with connecting lines and labeled in yellow, sites in commission are labeled in green, and legacy sites are labeled in red. From Paper II (Figure 1).

The sequence of Letters in this issue provides the full scope of the project and the conclusions drawn to date. Paper II opens with a description of the EHT array, the technical developments that enabled precursor detections, and the full range of observations reported here. Through the deployment of novel instrumentation at existing facilities, the collaboration created a new telescope with unique capabilities for black hole imaging. Paper III details the observations, data processing, calibration algorithms, and rigorous validation protocols for the final data products used for analysis. Paper IV gives the full process and approach to image reconstruction. The final images emerged after a rigorous evaluation of traditional imaging algorithms and new techniques tailored to the EHT instrument—alongside many months of testing the imaging algorithms through the analysis of synthetic data sets. Paper V uses newly assembled libraries of general relativistic magnetohydrodynamic (GRMHD) simulations and advanced ray-tracing to analyze the images and data in the context of black hole accretion and jet-launching. Paper VI employs model fits, comparison of simulations to data, and feature extraction from images to derive formal estimates of the lensed emission ring size and shape, black hole mass, and constraints on the nature of the black hole and the space-time surrounding it. Paper I is a concise summary.

Our image of the shadow confines the mass of M87 to within its photon orbit, providing the strongest case for the existence of supermassive black holes. These observations are consistent with Doppler brightening of relativistically moving plasma close to the black hole lensed around the photon orbit. They strengthen the fundamental connection between active galactic nuclei and central engines powered by accreting black holes through an entirely new approach. In the coming years, the EHT Collaboration will extend efforts to include full polarimetry, mapping of magnetic fields on horizon scales, investigations of time variability, and increased resolution through shorter wavelength observations.

In short, this work signals the development of a new field of research in astronomy and physics as we zero in on precision images of black holes on horizon scales. The prospects for sharpening our focus even further are excellent.

[First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole](#)

The Event Horizon Telescope Collaboration *et al.* 2019 *ApJL* **875** L1

[First M87 Event Horizon Telescope Results. II. Array and Instrumentation](#)

The Event Horizon Telescope Collaboration *et al.* 2019 *ApJL* **875** L2

[First M87 Event Horizon Telescope Results. III. Data Processing and Calibration](#)

The Event Horizon Telescope Collaboration *et al.* 2019 *ApJL* **875** L3

[First M87 Event Horizon Telescope Results. IV. Imaging the Central Supermassive Black Hole](#)

The Event Horizon Telescope Collaboration *et al.* 2019 *ApJL* **875** L4

[First M87 Event Horizon Telescope Results. V. Physical Origin of the Asymmetric Ring](#)

The Event Horizon Telescope Collaboration *et al.* 2019 *ApJL* **875** L5

[First M87 Event Horizon Telescope Results. VI. The Shadow and Mass of the Central Black Hole](#)

The Event Horizon Telescope Collaboration *et al.* 2019 *ApJL* **875** L6

First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole

The Event Horizon Telescope Collaboration, Kazunori Akiyama^{1,2,3,4} , Antxon Alberdi⁵ , Walter Alef⁶, Keiichi Asada⁷, Rebecca Azulay^{8,9,6} , Anne-Kathrin Baczko⁶ , David Ball¹⁰, Mislav Baloković^{4,11} , John Barrett² , Dan Bireley¹², Lindy Blackburn^{4,11} , Wilfred Boland¹³, Katherine L. Bouman^{4,11,14} , Geoffrey C. Bower¹⁵ , Michael Bremer¹⁶, Christiaan D. Brinkerink¹⁷ , Roger Brissenden^{4,11} , Silke Britzen⁶ , Avery E. Broderick^{18,19,20} , Dominique Brogiere¹⁶, Thomas Bronzwaer¹⁷, Do-Young Byun^{21,22} , John E. Carlstrom^{23,24,25,26}, Andrew Chael^{4,11} , Chi-kwan Chan^{10,27} , Shami Chatterjee²⁸ , Koushik Chatterjee²⁹, Ming-Tang Chen¹⁵, Yongjun Chen (陈永军)^{30,31}, Ilje Cho^{21,22} , Pierre Christian^{10,11} , John E. Conway³² , James M. Cordes²⁶, Geoffrey B. Crew² , Yuzhu Cui^{33,34} , Jordy Davelaar¹⁷ , Mariafelicia De Laurentis^{35,36,37} , Roger Deane^{38,39} , Jessica Dempsey¹² , Gregory Desvignes⁶ , Jason Dexter⁴⁰ , Sheperd S. Doeleman^{4,11} , Ralph P. Eatough⁶ , Helmo Falcke¹⁷ , Vincent L. Fish² , Ed Fomalont¹, Raquel Fraga-Escinas¹⁷ , William T. Freeman^{41,42}, Per Friberg¹², Christian M. Fromm³⁶, José L. Gómez⁵ , Peter Galison^{4,43,44} , Charles F. Gammie^{45,46} , Roberto García¹⁶, Olivier Gentaz¹⁶, Boris Georgiev^{19,20} , Ciriaco Goddi^{17,47}, Roman Gold³⁶ , Minfeng Gu (顾敏峰)^{30,48} , Mark Gurwell¹¹ , Kazuhiro Hada^{33,34} , Michael H. Hecht², Ronald Hesper⁴⁹ , Luis C. Ho (何子山)^{50,51} , Paul Ho⁷, Mareki Honma^{30,31} , Chih-Wei L. Huang⁷ , Lei Huang (黄磊)^{30,48}, David H. Hughes⁵², Shiro Ikeda^{3,53,54,55} , Makoto Inoue⁷, Sara Issaoun¹⁷ , David J. James^{4,11} , Buell T. Jannuzi¹⁰, Michael Janssen¹⁷ , Britton Jeter^{19,20} , Wu Jiang (江悟)³⁰ , Michael D. Johnson^{4,11} , Svetlana Jorstad^{56,57} , Taehyun Jung^{21,22} , Mansour Karami^{18,19} , Ramesh Karuppusamy⁸ , Tomohisa Kawashima³ , Garrett K. Keating⁴¹ , Mark Kettenis⁵⁸ , Jae-Young Kim⁶ , Junhan Kim¹⁰ , Jongsoo Kim²¹, Motoki Kino^{3,59} , Jun Yi Koay⁷ , Patrick M. Koch⁷ , Shoko Koyama⁷ , Michael Kramer⁶ , Carsten Kramer¹⁸ , Thomas P. Krichbaum⁶ , Cheng-Yu Kuo⁶⁰, Tod R. Lauer⁶¹ , Sang-Sung Lee²¹ , Yan-Rong Li (李彦荣)⁶² , Zhiyuan Li (李志远)^{63,64} , Michael Lindqvist³² , Kuo Liu⁶ , Elisabetta Liuzzo⁶⁵ , Wen-Ping Lo^{7,66}, Andrei P. Lobanov⁸, Laurent Loinard^{67,68} , Colin Lonsdale², Ru-Sen Lu (路如森)^{30,6} , Nicholas R. MacDonald⁶ , Jirong Mao (毛基荣)^{69,70,71} , Sera Markoff^{29,72} , Daniel P. Marrone¹⁰ , Alan P. Marscher⁵⁶ , Iván Martí-Vidal^{32,73} 

Satoki Matsushita⁷, Lynn D. Matthews² , Lia Medeiros^{10,74} , Karl M. Menten⁶ ,
 Yosuke Mizuno³⁶ , Izumi Mizuno¹² , James M. Moran^{4,11} , Kotaro Moriyama^{33,2} ,
 Monika Moscibrodzka¹⁷ , Cornelia Müller^{8,17} , Hiroshi Nagai^{3,34} , Neil M. Nagar⁷⁵ ,
 Masanori Nakamura⁷ , Ramesh Narayan^{4,11} , Gopal Narayanan⁷⁶, Iniyan Natarajan³⁹ ,
 Roberto Neri¹⁶, Chunchong Ni^{19,20} , Aristeidis Noutsos⁶ , Hiroki Okino^{33,77}, Héctor Olivares³⁸ ,
 Gisela N. Ortiz-León⁶ , Tomoaki Oyama³³, Feryal Özel¹⁰, Daniel C. M. Palumbo^{4,11} , Nimesh Patel¹¹,
 Ue-Li Pen^{18,78,79,80} , Dominic W. Pesce^{4,11} , Vincent Piétu¹⁶, Richard Plambeck⁸¹,
 Aleksandar PopStefanija⁷⁶, Oliver Porth^{29,36} , Ben Prather⁴⁵ , Jorge A. Preciado-López¹⁸ ,
 Dimitrios Psaltis¹⁰, Hung-Yi Pu¹⁸ , Venkatesh Ramakrishnan⁷⁵ , Ramprasad Rao¹⁵ ,
 Mark G. Rawlings¹², Alexander W. Raymond^{4,11} , Luciano Rezzolla³⁶ , Bart Ripperda³⁶ ,
 Freek Roelofs¹⁷ , Alan Rogers², Eduardo Ros⁸ , Mel Rose¹⁰ , Arash Roshanineshat¹⁰,
 Helge Rottmann⁶, Alan L. Roy⁶ , Chet Ruszczyk² , Benjamin R. Ryan^{80,83} , Kazi L. J. Rygl⁶⁵ ,
 Salvador Sánchez³⁴, David Sánchez-Argüelles^{52,85} , Mahito Sasada^{33,86} ,
 Tuomas Savolainen^{6,87,88} , F. Peter Schloerb⁷⁵, Karl-Friedrich Schuster¹⁵, Lijing Shao^{6,51} ,
 Zhiqiang Shen (沈志强)^{30,31} , Des Small⁵⁹ , Bong Won Sohn^{21,22,89} , Jason SooHoo² ,
 Fumie Tazaki³³ , Paul Tiede^{19,20} , Remo P. J. Tilanus^{17,47,90} , Michael Titus² ,
 Kenji Toma^{91,92} , Pablo Tome^{6,84} , Tyler Trent¹⁰, Sascha Trippe⁹³ , Shuichiro Tsuda³³,
 Ilse van Bommel⁵⁸ , Hulb Jan van Langevelde^{58,94} , Daniel R. van Rossum¹⁷ , Jan Wagner⁸,
 John Wardle⁹⁸ , Jonathan Weintraub^{4,11} , Norbert Wex⁶ , Robert Wharton⁶ ,
 Maciek Wielgus^{4,11} , George N. Wong⁴⁵ , Qingwen Wu (吴庆文)⁹⁶ , Ken Young¹¹ ,
 André Young¹⁷ , Ziri Younsi^{97,98} , Feng Yuan (袁峰)^{30,48,98} , Ye-Fei Yuan (袁业飞)⁹⁹,
 J. Anton Zensus⁶ , Guangyao Zhao²¹ , Shan-Shan Zhao^{17,83} , Ziyang Zhu⁴⁴,
 Juan-Carlos Algaba^{7,100} , Alexander Allardi¹⁰¹, Rodrigo Amestica¹⁰², Jacyndy Anczarski¹⁰³ ,
 Uwe Bach⁶ , Frederick K. Baganoff¹⁰⁴ , Christopher Beaudoin², Bradford A. Benson^{28,24} ,
 Ryan Berthold¹², Jay M. Blanchard^{75,58} , Ray Blundell¹¹, Sandra Bustamante¹⁰⁶, Roger Cappallo²,
 Edgar Castillo-Domínguez^{105,106}, Chih-Cheng Chang^{7,107}, Shu-Hao Chang⁷, Song-Chu Chang¹⁰⁷,
 Chung-Chen Chen⁷, Ryan Chilson¹⁵, Tim C. Chuter¹², Rodrigo Córdova Rosado^{4,11}, Iain M. Coulson¹² ,
 Thomas M. Crawford^{24,25} , Joseph Crowley¹⁰⁸, John David⁸⁴, Mark Derome², Matthew Dexter¹⁰⁹,
 Sven Dornbusch⁸, Kevin A. Duvovoi^{2,144}, Sergio A. Dzib⁶ , Andreas Eckart^{6,110} , Chris Eckart²,

Neal R. Erickson⁷⁶, Wendeline B. Everett¹¹¹ , Aaron Faber¹¹², Joseph R. Farah^{4,11,113} ,
 Vernon Fath⁷⁶, Thomas W. Folkers¹⁰, David C. Forbes¹⁰, Robert Freund¹⁰, Arturo I. Gómez-Ruiz^{105,106},
 David M. Gale¹⁰⁵, Feng Gao^{30,40}, Gertie Geertsema¹¹⁴, David A. Graham⁶, Christopher H. Greer¹⁰ ,
 Ronald Grosslein⁷⁶, Frédéric Gueth¹⁶, Daryl Haggard^{115,116,117} , Nils W. Halverson¹¹⁸ ,
 Chih-Chiang Han⁷, Kuo-Chang Han¹⁰⁷, Jinchi Hao¹⁰⁷, Yutaka Hasegawa⁷, Jason W. Henning^{23,119},
 Antonio Hernández-Gómez^{87,120} , Rubén Herrero-Illana¹²¹ , Stefan Heyminck⁸, Akihiko Hirota^{3,7},
 James Hoge¹², Yau-De Huang⁷, C. M. Violette Impellizzeri^{7,1}, Homn Jlang⁷, Atish Kamble^{4,11} ,
 Ryan Keisler²⁵ , Kimihiro Kimura⁷, Yusuke Kono³ , Derek Kubo¹²², John Kuroda¹²,
 Richard Lacasse¹⁰², Robert A. Laing¹²³, Erik M. Leitch²³ , Chao-Te Li⁷, Lupin C.-C. Lin^{7,124},
 Ching-Tang Liu¹⁰⁷, Kuan-Yu Liu⁷, Li-Ming Lu¹⁰⁷, Ralph G. Marson¹²⁵, Pierre L. Martin-Cocher⁷,
 Kyle D. Massingill¹⁰ , Callie Matulonis¹², Martin P. McColl¹⁰, Stephen R. McWhirter²,
 Hugo Messias^{121,126} , Zheng Meyer-Zhao^{7,127}, Daniel Michalik^{128,129} , Alfredo Montaña^{105,106},
 William Montgomerie¹², Matias Mora-Klein¹⁰², Dirk Muders⁶, Andrew Nadolski¹⁴⁸ , Santiago Navarro⁸⁴,
 Joseph Nelson¹⁰³ , Chi H. Nguyen^{10,130} , Hiroaki Nishioke⁷, Timothy Norton¹¹,
 Michael A. Nowak¹³¹ , George Nystrom¹⁵, Hideo Ogawa¹³², Peter Oshiro¹⁵, Tomoaki Oyama¹³³,
 Harriet Parsons¹² , Scott N. Paine¹¹ , Juan Peñalver⁸⁴, Neil M. Phillips^{121,126}, Michael Poirier²,
 Nicolas Pradel⁷, Rurik A. Primiani¹³⁴ , Philippe A. Raffin¹⁵, Alexandra S. Rahlin^{23,135} ,
 George Reiland¹⁰, Christopher Risacher¹⁶, Ignacio Ruiz⁶⁴, Alejandro F. Sáez-Madaín^{102,126},
 Remi Sassella¹⁶, Pim Schellart^{17,136} , Paul Shaw⁷, Kevin M. Silva¹², Hotaka Shiohawa¹¹ ,
 David R. Smith^{137,138} , William Snow¹⁵, Kamal Souccar⁷⁶, Don Sousa², T. K. Sridharan¹¹,
 Ranjani Srinivasan¹⁵, William Stahm¹², Anthony A. Stark¹¹ , Kyle Story¹³⁹, Sjoerd T. Timmer¹⁷ ,
 Laura Vertatschitsch^{11,134}, Craig Walther¹², Ta-Shun Wei⁷, Nathan Whitehorn¹⁴⁰ , Alan R. Whitney²,
 David P. Woody¹⁴¹, Jan G. A. Wouterloot¹² , Melvin Wright¹⁴² , Paul Yamaguchi¹¹ , Chen-Yu Yu⁷,
 Milagros Zeballos^{105,143}, Shuo Zhang¹⁰⁴ , and Lucy Ziurys¹⁰  — Hide full author list

Published 2019 April 10 • © 2019. The American Astronomical Society.

[The Astrophysical Journal Letters, Volume 875, Number 1](#)

[Focus on the First Event Horizon Telescope Results](#)



Article PDF



Article ePub

My friend, Carl

“No, it won’t be on the front page of every newspaper because the image is just too blurry”

The Boston Globe

THURSDAY, APRIL 11, 2019

Children's
hospital
prince
\$3.5m

royal promised
of child's care

z Kowalczyk
LOSS STAFF

Hospital has sued a member
nily for \$3.5 million, alleging
se to pay for the care of a very
one country.
Thursday in US District Court

One less mystery in our vast universe

Earthbound teams record
first image of black hole

By Brian MacQuarrie
GLOBE STAFF

Katherine Bouman had devoted years to the astonishing quest — to help capture the first image of a massive black hole in a distant galaxy, a void so dense no light can escape.

But when the mind-bending breakthrough finally came almost a year ago, the discovery had to stay a secret.

So, after the stunning image was revealed to the world Wednesday, Bouman's

excitement spilled out at what seemed the speed of light.

"We've been busting at the seams about what we've seen, but we had to keep our mouths shut," said Bouman, 29, a doctoral graduate of MIT who continued her studies at the Harvard-Smithsonian Center for Astrophysics.

What she and a large team of scientists from MIT, Harvard, and other universities had seen was the first-ever image of a cosmic black hole 55 million light-years away, a time-warping and light-twisting mystery of the universe whose existence Albert Einstein had hinted at a century ago.

BLACK HOLE, Page A12



EVENT HORIZON TELESCOPE COLLABORATION




The image of a black hole spawned celebrations.


Bernie Sanders
\$18.2 million

Warren

Middle of the pack total
cases concerns about
forgoing big donors







Katie Bouman


1 hr

Watching in disbelief as the first image I ever made of a black hole was in the process of being reconstructed.



37

6 Comments · 5 Shares



Share

View 2 more comments



Kaylan Balmanghelich This is amazing Katie! **Congratulations**

1h



Wardah Inam That is so cool!

1h



Shouni Puraglova You are so cute!

1h



Vikas Ramachandra very cool Katie!

47m



Adrian Dolce **Congrats** 🎉

36m



Mikl Rubincse **Congratulations!!**

17m

MIT CSAIL


3 years ago MIT grad student Katie Bouman led the creation of a new algorithm to produce the first-ever image of a black hole.

Today, that image was released.

More info: bit.ly/BHoleGuardian

2016 story: bit.ly/BlackHoleCSAIL

#EHTblackhole #EventHorizonTelescope




6:10 AM · 10 Apr 2019

MIT CSAIL
@MIT_CSAIL

MIT's largest research lab, the Computer Science & Artificial Intelligence Lab
[instagram.com/mit_csail/](https://www.instagram.com/mit_csail/) [facebook.com/mitcsail/](https://www.facebook.com/mitcsail/) [youtube.com/mitcsail/](https://www.youtube.com/channel/UCv3p0D8333333333333333) [linkedin.com/company/mit-csail/](https://www.linkedin.com/company/mit-csail/) [github.com/mit-csail/](https://www.github.com/mit-csail/) [mit.edu/csail/](https://www.mit.edu/csail/)

Cambridge, MA
csail@mit.edu
Joined October 2009



MIT CSAIL

@MIT_CSAIL

MIT's largest research lab, the Computer Science & Artificial Intelligence Lab
instagram.com/mit_csail/ @ai @ml #bigdata
that trip. twireless

Cambridge, MA

csail.mit.edu

Joined October 2009





MIT CSAIL

@MIT_CSAIL

Follow

Here's the moment when the first black hole image was processed, from the eyes of researcher Katie Bouman. #EHTBlackHole #BlackHoleDay #BlackHole (v/@dfbarajas)



9:51 AM - 10 Apr 2019

17,550 Retweets 58,160 Likes



579 18K 58K



9400

Congresswoman for NY-14 (Bronx + Queens). In a modern, moral, + wealthy society, no American should be too poor to live. ~~NO~~ % People-Funded, no corporate PAC \$

© Bronx + Queens, NYC

Joined April 2010



9AOC

Follow



Here's to [#WomenInSTEM!](#)



Congratulations to Katie Beuman to whom we owe the first photograph of a black hole ever. Not seeing her name circulate nearly enough in the press....

1:41 PM - 10 Apr 2019

23,504 Retweets 101,643 Likes



1.0K 24K 102K



Replying to SAOC

And "she" did by hand all by herself right? All with the help of her vagina right? Give me a break. Scientific and technological progress has no sex.



Nancy Pelosi ✓

@SpeakerPelosi



Congratulations to Dr. Katie Bouman, who developed the algorithm which captured the first ever image of a black hole! You are an inspiration to all Americans and especially to young women and girls with STEAM dreams! [cnn.com/2019/04/10/us/...](https://www.cnn.com/2019/04/10/us/...)

♡ 40.3K 6:46 PM - Apr 11, 2019



That image of a black hole you saw everywh...

The effort wouldn't have succeeded without Katie Bouman, who developed a crucial algorithm and [cnn.com](https://www.cnn.com)



Barbie ✓

@Barbie




Congratulations Katie Bouman on this remarkable accomplishment! Thank you for leading by example and encouraging girls to push the boundaries of science. 🧠
[#YouCanBeAnything](#) [#MoreRoleModels](#)

AJ+ ✓ @ajplus

Replying to @ajplus

This is Dr. Katie Bouman. She's the computer scientist behind the first-ever image of a black hole.

She developed the algorithm that turned telescopic data into the historic photo we see today.



♡ 550 6:46 PM - Apr 10, 2019



🗨 150 people are talking about this



Read more:



MIT CSAIL ✓

@MIT_CSAIL

Follow



We at [@MIT_CSAIL](#) are so proud of the role our alum Dr. Katie Bouman played in the development of the first-ever picture of a black hole. She's been psyched about all the [#blackhole](#) interest & just wanted to clarify a few things. (1/7)

11:49 AM · 12 Apr 2019

128 Retweets 434 Likes



26

128

434



MIT CSAIL ✓ @MIT_CSAIL · Apr 12



In our first tweet about this, we linked to a 2016 story about an algorithm she led the development of while at CSAIL. That algorithm was intended to take a picture of a black hole, but didn't create the final image. (cont.)(2/7)

4

11

81



MIT CSAIL ✓ @MIT_CSAIL · Apr 12



It inspired image validation procedures in the final paper, and the EHT team together developed new methods that were used in reconstructing the black hole image.(3/7)

2

9

79

© 2019 Twitter · About · Help

Privacy policy · Cookies

How Katie Bouman Accidentally Became the Face of the Black Hole Project

By Sarah Mervosh

April 11, 2019

[Leer en español](#)

As the first-ever picture of a black hole [was unveiled](#) this week, another image began making its way around the internet: a [photo of a young scientist](#), clasping her hands over her face and reacting with glee to an image of an orange ring of light, circling a deep, dark abyss.

It was a photo too good not to share. The scientist, Katie Bouman, a postdoctoral fellow who contributed to the project, became an instant hero for women and girls in STEM, a welcome symbol in a world hungry for representation.

Public figures from Washington to Hollywood learned her name. And [some advocates](#), familiar with how history can

New York Times

The Atlantic


☰

SCIENCE

The Dark Saga of Katie Bouman

How a young scientist got sucked into the black hole of the internet

MARINA KOREN 1:16 PM ET



DESIGN / GETTY / EHT / NATIONAL SCIENCE FOUNDATION VIA REUTERS / THE ATLANTIC

Two photos—one long anticipated, the other a



People also view

Katie Bouman: Black hole colleagues dismiss attacks over hero scientist's role in creating famous image



The Atlantic



Katie Bouman is with Sara Issaoun and 6 others.


April 10 at 7:47 PM · 🌐

I'm so excited that we finally get to share what we have been working on for the past year! The image shown today is the combination of images produced by multiple methods. No one algorithm or person made this image, it required the amazing talent of a team of scientists from around the globe and years of hard work to develop the instrument, data processing, imaging methods, and analysis techniques that were necessary to pull off this seemingly impossible feat. It has been truly an honor, and I am so lucky to have had the opportunity to work with you all.



👍❤️😱 71K

8.8K Comments 9.6K Shares



Katie speaking
at MIT



Katie speaking
at MIT



Katie speaking
at Stanford



Katie's audience
at Stanford



Katie speaking at CSAIL's annual gala at ICA, April 27, 2019



Franny, Bill, Katie, Joe at ICA



next up: SgrA*

- The EHT team is processing the data from the black hole at the center of our galaxy
- Much less massive (4M vs 6.5B solar masses), and therefore faster dynamics than M87*
- They're working on processing the SgrA* data now...